

Aspects of T-system and Y-system

Atsuo Kuniba (Univ. Tokyo)

20 May 2010

Komaba Particle Theory Group Weekly Seminar

Contents

- What are T-systems and Y-systems?
- Restriction and periodicity
- Quivers and Cluster algebra formulation
- Dilogarithm identity

What are T-systems and Y-systems?

Systems of difference equations among commuting variables

$$T_m^{(a)}(u) \quad \text{and} \quad Y_m^{(a)}(u)$$


related to root system.

$$a \in \{\text{nodes of Dynkin diagram of } \mathfrak{g}\}$$
$$(\mathfrak{g} = A_n, B_n, C_n, D_n, E_6, E_7, E_8, F_4, G_2)$$

$$m \in \mathbb{Z}_{\geq 1}$$

$$u \in \mathbb{C} \quad (\text{spectral parameter})$$

$$t_a := |\text{long root}|^2 / |\alpha_a|^2 \quad (= 1 \text{ for ADE})$$



$$T_m^{(a)}\left(u - \frac{1}{t_a}\right) T_m^{(a)}\left(u + \frac{1}{t_a}\right) = T_{m-1}^{(a)}(u) T_{m+1}^{(a)}(u) + \text{product of } T\text{'s,}$$

$$Y_m^{(a)}\left(u - \frac{1}{t_a}\right) Y_m^{(a)}\left(u + \frac{1}{t_a}\right) = \frac{\text{product of } (1 + Y)\text{'s}}{(1 + Y_{m-1}^{(a)}(u))^{-1} (1 + Y_{m+1}^{(a)}(u))^{-1}}.$$

Structure of products in the RHS is dependent on $m \bmod t_a \mathbb{Z}$.

$\mathfrak{g} = A_n, D_n, E_n$ case

$C = (C_{ab})_{1 \leq a, b \leq n}$: Cartan matrix

T-system

$$T_m^{(a)}(u-1)T_m^{(a)}(u+1) = T_{m-1}^{(a)}(u)T_{m+1}^{(a)}(u) + \prod_{b: C_{ab}=-1} T_m^{(b)}(u)$$

Example A_n

$$T_m^{(a)}(u-1)T_m^{(a)}(u+1) = T_{m-1}^{(a)}(u)T_{m+1}^{(a)}(u) + T_m^{(a-1)}(u)T_m^{(a+1)}(u)$$
$$(T_0^{(a)}(u) = T_m^{(0)}(u) = T_m^{(n+1)}(u) = 1.)$$

A version of Hirota-Miwa or Toda-field equation
on discrete space-time.

B_n

$$\begin{aligned}T_m^{(a)}(u-1)T_m^{(a)}(u+1) &= T_{m-1}^{(a)}(u)T_{m+1}^{(a)}(u) + T_m^{(a-1)}(u)T_m^{(a+1)}(u) \quad (1 \leq a \leq n-2), \\T_m^{(n-1)}(u-1)T_m^{(n-1)}(u+1) &= T_{m-1}^{(n-1)}(u)T_{m+1}^{(n-1)}(u) + T_m^{(n-2)}(u)T_{2m}^{(n)}(u), \\T_{2m}^{(n)}(u-\frac{1}{2})T_{2m}^{(n)}(u+\frac{1}{2}) &= T_{2m-1}^{(n)}(u)T_{2m+1}^{(n)}(u) + T_m^{(n-1)}(u-\frac{1}{2})T_m^{(n-1)}(u+\frac{1}{2}), \\T_{2m+1}^{(n)}(u-\frac{1}{2})T_{2m+1}^{(n)}(u+\frac{1}{2}) &= T_{2m}^{(n)}(u)T_{2m+2}^{(n)}(u) + T_m^{(n-1)}(u)T_{m+1}^{(n-1)}(u).\end{aligned}$$

C_n

$$\begin{aligned}T_m^{(a)}(u-\frac{1}{2})T_m^{(a)}(u+\frac{1}{2}) &= T_{m-1}^{(a)}(u)T_{m+1}^{(a)}(u) + T_m^{(a-1)}(u)T_m^{(a+1)}(u) \quad (1 \leq a \leq n-2), \\T_{2m}^{(n-1)}(u-\frac{1}{2})T_{2m}^{(n-1)}(u+\frac{1}{2}) &= T_{2m-1}^{(n-1)}(u)T_{2m+1}^{(n-1)}(u) + T_{2m}^{(n-2)}(u)T_m^{(n)}(u-\frac{1}{2})T_m^{(n)}(u+\frac{1}{2}), \\T_{2m+1}^{(n-1)}(u-\frac{1}{2})T_{2m+1}^{(n-1)}(u+\frac{1}{2}) &= T_{2m}^{(n-1)}(u)T_{2m+2}^{(n-1)}(u) + T_{2m+1}^{(n-2)}(u)T_m^{(n)}(u)T_{m+1}^{(n)}(u), \\T_m^{(n)}(u-1)T_m^{(n)}(u+1) &= T_{m-1}^{(n)}(u)T_{m+1}^{(n)}(u) + T_{2m}^{(n-1)}(u).\end{aligned}$$

F_4

$$T_m^{(1)}(u-1)T_m^{(1)}(u+1) = T_{m-1}^{(1)}(u)T_{m+1}^{(1)}(u) + T_m^{(2)}(u),$$

$$T_m^{(2)}(u-1)T_m^{(2)}(u+1) = T_{m-1}^{(2)}(u)T_{m+1}^{(2)}(u) + T_m^{(1)}(u)T_{2m}^{(3)}(u),$$

$$T_{2m}^{(3)}(u - \frac{1}{2})T_{2m}^{(3)}(u + \frac{1}{2}) = T_{2m-1}^{(3)}(u)T_{2m+1}^{(3)}(u) + T_m^{(2)}(u - \frac{1}{2})T_m^{(2)}(u + \frac{1}{2})T_{2m}^{(4)}(u),$$

$$T_{2m+1}^{(3)}(u - \frac{1}{2})T_{2m+1}^{(3)}(u + \frac{1}{2}) = T_{2m}^{(3)}(u)T_{2m+2}^{(3)}(u) + T_m^{(2)}(u)T_{m+1}^{(2)}(u)T_{2m+1}^{(4)}(u),$$

$$T_m^{(4)}(u - \frac{1}{2})T_m^{(4)}(u + \frac{1}{2}) = T_{m-1}^{(4)}(u)T_{m+1}^{(4)}(u) + T_m^{(3)}(u).$$

G_2

$$T_m^{(1)}(u-1)T_m^{(1)}(u+1) = T_{m-1}^{(1)}(u)T_{m+1}^{(1)}(u) + T_{3m}^{(2)}(u),$$

$$T_{3m}^{(2)}(u - \frac{1}{3})T_{3m}^{(2)}(u + \frac{1}{3}) = T_{3m-1}^{(2)}(u)T_{3m+1}^{(2)}(u) + T_m^{(1)}(u - \frac{2}{3})T_m^{(1)}(u)T_m^{(1)}(u + \frac{2}{3}),$$

$$T_{3m+1}^{(2)}(u - \frac{1}{3})T_{3m+1}^{(2)}(u + \frac{1}{3}) = T_{3m}^{(2)}(u)T_{3m+2}^{(2)}(u) + T_m^{(1)}(u - \frac{1}{3})T_m^{(1)}(u + \frac{1}{3})T_{m+1}^{(1)}(u),$$

$$T_{3m+2}^{(2)}(u - \frac{1}{3})T_{3m+2}^{(2)}(u + \frac{1}{3}) = T_{3m+1}^{(2)}(u)T_{3m+3}^{(2)}(u) + T_m^{(1)}(u)T_{m+1}^{(1)}(u - \frac{1}{3})T_{m+1}^{(1)}(u + \frac{1}{3}).$$

Origin of T-system: $T_m^{(a)}(u)$ stands for

Phys: commuting transfer matrices in
Yang-Baxter solvable lattice models.

$$T_m^{(a)}(u) = \text{Tr}_{W_m^{(a)}(u)} \left(\begin{array}{c} u \\ \hline | \quad | \quad | \quad \cdots \quad | \end{array} \right), \quad [T_m^{(a)}(u), T_{m'}^{(a')}(u')] = 0.$$

Math: q -characters of Kirillov-Reshetikhin modules $W_m^{(a)}(u)$ of
quantum affine algebra $U_q(\hat{\mathfrak{g}})$.

$$\text{For } A_n \quad W_m^{(a)}(u) \simeq \widehat{\square}^m a$$

$$0 \rightarrow W_m^{(a-1)}(u) \otimes W_m^{(a+1)}(u) \rightarrow W_m^{(a)}(u-1) \otimes W_m^{(a)}(u+1) \rightarrow W_{m-1}^{(a)}(u) \otimes W_{m+1}^{(a)}(u) \rightarrow 0$$

Proposed in the former context by K-Nakanishi-Suzuki (1994).

Proved in the latter context by Nakajima for ADE (2003)
and Hernandez for $\forall \mathfrak{g}$ (2006).

Y-system

$$\mathfrak{g} = A_n, D_n, E_n \text{ case} \quad (Y_0^{(a)}(\mathbf{u})^{-1} = 0)$$

$$Y_m^{(a)}(\mathbf{u} - 1)Y_m^{(a)}(\mathbf{u} + 1) = \frac{\prod_{b: C_{ab}=-1} (1 + Y_m^{(b)}(\mathbf{u}))}{(1 + Y_{m-1}^{(a)}(\mathbf{u})^{-1})(1 + Y_{m+1}^{(a)}(\mathbf{u})^{-1})}$$

B_n

$$Y_m^{(a)}(\mathbf{u} - 1)Y_m^{(a)}(\mathbf{u} + 1) = \frac{(1 + Y_m^{(a-1)}(\mathbf{u}))(1 + Y_m^{(a+1)}(\mathbf{u}))}{(1 + Y_{m-1}^{(a)}(\mathbf{u})^{-1})(1 + Y_{m+1}^{(a)}(\mathbf{u})^{-1})} \quad (1 \leq a \leq n - 2),$$

$$\begin{aligned} & Y_m^{(n-1)}(\mathbf{u} - 1)Y_m^{(n-1)}(\mathbf{u} + 1) \\ &= \frac{(1 + Y_m^{(n-2)}(\mathbf{u}))(1 + Y_{2m}^{(n)}(\mathbf{u} + \frac{1}{2}))(1 + Y_{2m}^{(n)}(\mathbf{u} - \frac{1}{2}))(1 + Y_{2m-1}^{(n)}(\mathbf{u}))(1 + Y_{2m+1}^{(n)}(\mathbf{u}))}{(1 + Y_{m-1}^{(n-1)}(\mathbf{u})^{-1})(1 + Y_{m+1}^{(n-1)}(\mathbf{u})^{-1})}, \end{aligned}$$

$$Y_{2m}^{(n)}(\mathbf{u} - \frac{1}{2})Y_{2m}^{(n)}(\mathbf{u} + \frac{1}{2}) = \frac{1 + Y_m^{(n-1)}(\mathbf{u})}{(1 + Y_{2m-1}^{(n)}(\mathbf{u})^{-1})(1 + Y_{2m+1}^{(n)}(\mathbf{u})^{-1})},$$

$$Y_{2m+1}^{(n)}(\mathbf{u} - \frac{1}{2})Y_{2m+1}^{(n)}(\mathbf{u} + \frac{1}{2}) = \frac{1}{(1 + Y_{2m}^{(n)}(\mathbf{u})^{-1})(1 + Y_{2m+2}^{(n)}(\mathbf{u})^{-1})}.$$

C_n

$$Y_m^{(a)}\left(u - \frac{1}{2}\right)Y_m^{(a)}\left(u + \frac{1}{2}\right) = \frac{(1 + Y_m^{(a-1)}(u))(1 + Y_m^{(a+1)}(u))}{(1 + Y_{m-1}^{(a)}(u)^{-1})(1 + Y_{m+1}^{(a)}(u)^{-1})} \quad (1 \leq a \leq n - 2),$$

$$Y_{2m}^{(n-1)}\left(u - \frac{1}{2}\right)Y_{2m}^{(n-1)}\left(u + \frac{1}{2}\right) = \frac{(1 + Y_{2m}^{(n-2)}(u))(1 + Y_m^{(n)}(u))}{(1 + Y_{2m-1}^{(n-1)}(u)^{-1})(1 + Y_{2m+1}^{(n-1)}(u)^{-1})},$$

$$Y_{2m+1}^{(n-1)}\left(u - \frac{1}{2}\right)Y_{2m+1}^{(n-1)}\left(u + \frac{1}{2}\right) = \frac{1 + Y_{2m+1}^{(n-2)}(u)}{(1 + Y_{2m}^{(n)}(u)^{-1})(1 + Y_{2m+2}^{(n)}(u)^{-1})},$$

$$\begin{aligned} & Y_m^{(n)}(u - 1)Y_m^{(n)}(u + 1) \\ &= \frac{(1 + Y_{2m}^{(n-1)}\left(u + \frac{1}{2}\right))(1 + Y_{2m}^{(n-1)}\left(u - \frac{1}{2}\right))(1 + Y_{2m-1}^{(n-1)}(u))(1 + Y_{2m+1}^{(n-1)}(u))}{(1 + Y_{m-1}^{(n)}(u)^{-1})(1 + Y_{m+1}^{(n)}(u)^{-1})}. \end{aligned}$$

F_4

$$\begin{aligned} Y_m^{(1)}(u-1)Y_m^{(1)}(u+1) &= \frac{1 + Y_m^{(2)}(u)}{(1 + Y_{m-1}^{(1)}(u)^{-1})(1 + Y_{m+1}^{(1)}(u)^{-1})}, \\ Y_m^{(2)}(u-1)Y_m^{(2)}(u+1) &= \frac{(1 + Y_m^{(1)}(u))(1 + Y_{2m}^{(3)}(u - \frac{1}{2}))(1 + Y_{2m}^{(3)}(u + \frac{1}{2}))(1 + Y_{2m-1}^{(3)}(u))(1 + Y_{2m+1}^{(3)}(u))}{(1 + Y_{m-1}^{(2)}(u)^{-1})(1 + Y_{m+1}^{(2)}(u)^{-1})}, \\ Y_{2m}^{(3)}(u - \frac{1}{2})Y_{2m}^{(3)}(u + \frac{1}{2}) &= \frac{(1 + Y_m^{(2)}(u))(1 + Y_{2m}^{(4)}(u))}{(1 + Y_{2m-1}^{(3)}(u)^{-1})(1 + Y_{2m+1}^{(3)}(u)^{-1})}, \\ Y_{2m+1}^{(3)}(u - \frac{1}{2})Y_{2m+1}^{(3)}(u + \frac{1}{2}) &= \frac{1 + Y_{2m+1}^{(4)}(u)}{(1 + Y_{2m}^{(3)}(u)^{-1})(1 + Y_{2m+2}^{(3)}(u)^{-1})}, \\ Y_m^{(4)}(u - \frac{1}{2})Y_m^{(4)}(u + \frac{1}{2}) &= \frac{1 + Y_m^{(3)}(u)}{(1 + Y_{m-1}^{(4)}(u)^{-1})(1 + Y_{m+1}^{(4)}(u)^{-1})}. \end{aligned}$$

G_2

$$\begin{aligned} Y_m^{(1)}(u-1)Y_m^{(1)}(u+1) &= (1 + Y_{3m}^{(2)}(u - \frac{2}{3}))(1 + Y_{3m}^{(2)}(u))(1 + Y_{3m}^{(2)}(u + \frac{2}{3})) \\ &\quad \times (1 + Y_{3m-1}^{(2)}(u - \frac{1}{3}))(1 + Y_{3m-1}^{(2)}(u + \frac{1}{3})) \\ &\quad \times (1 + Y_{3m+1}^{(2)}(u - \frac{1}{3}))(1 + Y_{3m+1}^{(2)}(u + \frac{1}{3})) \\ &\quad \times (1 + Y_{3m-2}^{(2)}(u))(1 + Y_{3m+2}^{(2)}(u)) \\ &\quad \times \left((1 + Y_{m-1}^{(1)}(u)^{-1})(1 + Y_{m+1}^{(1)}(u)^{-1}) \right)^{-1} \end{aligned}$$

$$Y_{3m}^{(2)}(u - \frac{1}{3})Y_{3m}^{(2)}(u + \frac{1}{3}) = \frac{1 + Y_m^{(1)}(u)}{(1 + Y_{3m-1}^{(2)}(u)^{-1})(1 + Y_{3m+1}^{(2)}(u)^{-1})},$$

$$Y_{3m+1}^{(2)}(u - \frac{1}{3})Y_{3m+1}^{(2)}(u + \frac{1}{3}) = \frac{1}{(1 + Y_{3m}^{(2)}(u)^{-1})(1 + Y_{3m+2}^{(2)}(u)^{-1})},$$

$$Y_{3m+2}^{(2)}(u - \frac{1}{3})Y_{3m+2}^{(2)}(u + \frac{1}{3}) = \frac{1}{(1 + Y_{3m+1}^{(2)}(u)^{-1})(1 + Y_{3m+3}^{(2)}(u)^{-1})}.$$

Y-system is an algebraic form of thermodynamic Bethe ansatz equation of type \mathfrak{g} under string hypothesis.

$Y_m^{(a)}(u) \sim$ Boltzmann factor of string/hole excitation
with color a , length m , rapidity u .

A_1 example: $(Y_m(u) = Y_m^{(1)}(u)^{-1})$

$$\log Y_m(u) = \text{known fcn.} + \int_{-\infty}^{\infty} \frac{\log(1 + Y_{m-1}(v))(1 + Y_{m+1}(v))}{4 \cosh \frac{\pi(u-v)}{2}} dv$$

$$\rightsquigarrow Y_m(u - i)Y_m(u + i) = (1 + Y_{m-1}(u))(1 + Y_{m+1}(u)).$$

Y-system was proposed by

ADE: Al. Zamolodchikov (1991), Ravanini-Tateo-Valleriani (1993).

$\forall \mathfrak{g}$: K-Nakanishi (1992).

Relation of T and Y-systems

A_1 example

$$\begin{aligned} Y_m(u-1)Y_m(u+1) &= (1 + Y_{m-1}(u))(1 + Y_{m+1}(u)), \\ T_m(u-1)T_m(u+1) &= T_{m-1}(u)T_{m+1}(u) + 1. \end{aligned}$$

Formally setting $Y_m(u) = T_{m-1}(u)T_{m+1}(u)$,

$$\begin{aligned} Y_m(u-1)Y_m(u+1) &= T_{m-1}(u-1)T_{m+1}(u-1)T_{m-1}(u+1)T_{m+1}(u+1) \\ &= T_{m+1}(u-1)T_{m+1}(u+1)T_{m-1}(u-1)T_{m-1}(u+1) \\ &= (T_{m+2}(u)T_m(u) + 1)(T_{m-2}(u)T_m(u) + 1) \\ &= (Y_{m+1}(u) + 1)(Y_{m-1}(u) + 1). \end{aligned}$$

Similarly for $\forall \mathfrak{g}$,

T-system solves Y-system.

(Yet to be understood why.)

Restriction and Periodicity

Introduce $\ell \in \mathbb{Z}_{\geq 2}$ called **level**.

Level ℓ restricted T and Y-system are those closing among

$$T_m^{(a)}(u) \text{ and } Y_m^{(a)}(u) \text{ with } 1 \leq m \leq t_a \ell - 1,$$

obtained respectively by imposing

$$T_{t_a \ell}^{(a)}(u) = 1 \text{ and } Y_{t_a \ell}^{(a)}(u)^{-1} = 0.$$

C_2 example. $(t_1, t_2) = (2, 1)$

$$\begin{aligned} T_{2m+1}^{(1)}\left(u - \frac{1}{2}\right)T_{2m+1}^{(1)}\left(u + \frac{1}{2}\right) &= T_{2m}^{(1)}(u)T_{2m+2}^{(1)}(u) + T_m^{(2)}(u)T_{m+1}^{(2)}(u), \\ T_{2m}^{(1)}\left(u - \frac{1}{2}\right)T_{2m}^{(1)}\left(u + \frac{1}{2}\right) &= T_{2m-1}^{(1)}(u)T_{2m+1}^{(1)}(u) + T_m^{(2)}\left(u - \frac{1}{2}\right)T_m^{(2)}\left(u + \frac{1}{2}\right), \\ T_m^{(2)}(u-1)T_m^{(2)}(u+1) &= T_{m-1}^{(2)}(u)T_{m+1}^{(2)}(u) + T_{2m}^{(1)}(u). \end{aligned}$$

Level 2 restriction: $T_4^{(1)}(u) = T_2^{(2)}(u) = 1$. (b.c. $T_0^{(a)}(u) = 1$)

$$\begin{aligned} T_1^{(1)}\left(u - \frac{1}{2}\right)T_1^{(1)}\left(u + \frac{1}{2}\right) &= T_2^{(1)}(u) + T_1^{(2)}(u), \\ T_2^{(1)}\left(u - \frac{1}{2}\right)T_2^{(1)}\left(u + \frac{1}{2}\right) &= T_1^{(1)}(u)T_3^{(1)}(u) + T_1^{(1)}\left(u - \frac{1}{2}\right)T_1^{(1)}\left(u + \frac{1}{2}\right), \\ T_3^{(1)}\left(u - \frac{1}{2}\right)T_3^{(1)}\left(u + \frac{1}{2}\right) &= T_2^{(1)}(u) + T_1^{(2)}(u), \\ T_1^{(2)}(u-1)T_1^{(2)}(u+1) &= 1 + T_2^{(1)}(u), \end{aligned}$$

which closes among $T_1^{(1)}(u), T_2^{(1)}(u), T_3^{(1)}(u), T_1^{(2)}(u)$.


Restricted T and Y-systems \dots evolution eqs. in the u direction.

Periodicity conjecture

Level ℓ restricted T -system and Y -system obey

$$T_m^{(a)}(u + h^\vee + \ell) = T_{t_a \ell - m}^{(\omega(a))}(u) \quad \text{and} \quad Y_m^{(a)}(u + h^\vee + \ell) = Y_{t_a \ell - m}^{(\omega(a))}(u).$$

ω is an involution whose only non-trivial cases are



$h^\vee =$ dual Coxeter number

\mathfrak{g}	A_n	B_n	C_n	D_n	E_6	E_7	E_8	F_4	G_2
h^\vee	$n+1$	$2n-1$	$n+1$	$2n-2$	12	18	30	9	4

(Full periodicity: $T_m^{(a)}(u + 2(h^\vee + \ell)) = T_m^{(a)}(u)$ and same for $Y_m^{(a)}(u)$.)

Example: $(A_2, 2)$

Write $T^{(a)}(u) = T_1^{(a)}(u)$.

$$T^{(1)}(u-1)T^{(1)}(u+1) = 1 + T^{(2)}(u),$$

$$T^{(2)}(u-1)T^{(2)}(u+1) = 1 + T^{(1)}(u).$$

Periodicity reads

$$T^{(1)}(u+5) = T^{(2)}(u), \quad T^{(2)}(u+5) = T^{(1)}(u).$$

$$T^{(1)}(0) = a$$

$$T^{(2)}(1) = b$$

$$T^{(1)}(2) = \frac{1 + T^{(2)}(1)}{T^{(1)}(0)} = \frac{1 + b}{a}$$

$$T^{(2)}(3) = \frac{1 + T^{(1)}(2)}{T^{(2)}(1)} = \frac{1 + \frac{1+b}{a}}{b} = \frac{1 + a + b}{ab}$$

$$T^{(1)}(4) = \frac{1 + T^{(2)}(3)}{T^{(1)}(2)} = \frac{1 + \frac{1+a+b}{ab}}{\frac{1+b}{a}} = \frac{1 + a}{b}$$

$$T^{(2)}(5) = \frac{1 + T^{(1)}(4)}{T^{(2)}(3)} = \frac{1 + \frac{1+a}{b}}{\frac{1+a+b}{ab}} = a = T^{(1)}(0)$$

$$T^{(1)}(6) = \frac{1 + T^{(2)}(5)}{T^{(1)}(4)} = \frac{1 + a}{\frac{1+a}{b}} = b = T^{(2)}(1)$$

0{10, 30, 50, 70}

$$(E_8, 2) : \{T_1^{(1)}(u), T_1^{(3)}(u), T_1^{(5)}(u), T_1^{(7)}(u)\}_{u=0}^{32}$$

$$2 \left\{ \frac{11}{5}, \frac{431}{15}, \frac{101291}{25}, \frac{31}{35} \right\}$$

$$4 \left\{ \frac{83}{45}, \frac{69696833}{230625}, \frac{45718438593497}{22157296875}, \frac{103041}{1525} \right\}$$

$$6 \left\{ \frac{102041}{1025}, \frac{8821291833971}{66471890625}, \frac{360342463107797294639}{34634624677734375}, \frac{14562107}{415125} \right\}$$

$$8 \left\{ \frac{1061807}{1441125}, \frac{527621002287915653}{153931665234375}, \frac{144652414821069001465529527}{6161870815433349609375}, \frac{2176297573}{492384375} \right\}$$

$$10 \left\{ \frac{15241182}{312625}, \frac{17418588023516754184}{133590695185546875}, \frac{65852952390687824418240896525206}{1926354863674850921630859375}, \frac{32206227374}{211021875} \right\}$$

$$12 \left\{ \frac{23381761}{6226875}, \frac{4439405789261107709041}{9128697504345703125}, \frac{255396681651083275452908699280166448}{8813073501312442966461181640625}, \frac{6587423634821}{129778453125} \right\}$$

$$14 \left\{ \frac{289412993}{98476875}, \frac{2401172003278457388295019}{2875539713868896484375}, \frac{113421595121251725116844505024021577713}{5420040203307152424373626708984375}, \frac{8472179120234}{2252658515625} \right\}$$

$$16 \left\{ \frac{391949128}{4689375}, \frac{7397263161797774132227049}{58469307515334228515625}, \frac{1290705517162033306270461619591091257193}{569104221347251004559230804443359375}, \frac{14335608965944}{129778453125} \right\}$$

$$18 \left\{ \frac{66998956}{126613125}, \frac{210714979567782348600241}{928084246275146484375}, \frac{172470738440320575058431884494833913663}{113820844269450200911846160888671875}, \frac{74693044181731}{13626737578125} \right\}$$

$$20 \left\{ \frac{11232037}{1563125}, \frac{1576259942401957743647}{246474832617333984375}, \frac{2104768617341673326572332456823959011}{2529352094876671131374359130859375}, \frac{1211696207719}{450531703125} \right\}$$

$$22 \left\{ \frac{3077201}{1245375}, \frac{115401582866988182927}{4260058835361328125}, \frac{23354104411061828987973549647671}{2467660580367484030609130859375}, \frac{175786811543}{2883965625} \right\}$$

$$24 \left\{ \frac{4476646}{2188375}, \frac{47183886310350193}{12468464883984375}, \frac{35939455246726991953433003}{1712315434377645263671875}, \frac{991662341}{13294378125} \right\}$$

$$26 \left\{ \frac{7058}{4575}, \frac{1842216632119}{879609515625}, \frac{131289331831932106159}{115021588554755859375}, \frac{7222892}{312625} \right\}$$

$$28 \left\{ \frac{1181}{615}, \frac{61893029}{42204375}, \frac{156275914764469}{18471799828125}, \frac{46522}{27675} \right\}$$

$$30 \left\{ \frac{23}{15}, \frac{32333}{1845}, \frac{4966808}{187575}, \frac{1781}{2135} \right\}$$

32{10, 30, 50, 70}

Periodicity of Y-system for (\mathfrak{g}, ℓ) was proposed:

Al. Zamolodchikov (1991) (ADE, 2),

Ravanini-Tateo-Valleriani (1993) (ADE, ℓ),

K-Nakanishi-Suzuki (1994) (\mathfrak{g}, ℓ) .

Many cases proved in

$(A_n, 2)$: Frenkel-Szenes (1995), Gliozzi-Tateo (1996),

(A_n, ℓ) : Volkov, Henriques (2007),

(ADE, 2): Fomin-Zelevinsky (2003) [Cluster algebra](#),

(ADE, ℓ): Keller (arXiv:0807.1960) [Cluster algebra](#)/[category](#).

Periodicity of T-system:

Proposed and partially proved in Inoue-Iyama-K-Nakanishi-Suzuki (2008),

(A_n, ℓ) case: proof also contained in Henriques (2007).

[Theorem. \[Inoue-Iyama-Keller-K-Nakanishi 2010\]](#)

[Periodicity conjecture of T and Y-systems is true for \$\forall\(\mathfrak{g}, \ell\)\$.](#)

Quivers and Cluster algebra formulation

Q : quiver (finite oriented graph without loop  and 2-cycle )

$I = \{1, \dots, N\}$: vertex set, $x = (x_1, \dots, x_N)$: I -tuple of variables

x_i : **cluster variable**, (Q, x) : **seed**.

Cluster algebra \mathcal{A}_Q is defined by (i)–(iv). Fomin-Zelevinsky (2002)

(i) Start from the (initial) seed (Q, x) as above.


(ii) For each $k \in I$, define another seed (R, y)
by $(R, y) = \mu_k(Q, x)$ (**‘mutation’ at k** , def. next page).

(iii) Iterate mutations for every new seed at every k ,
and collect all (possibly infinite) seeds.

(iv) $\mathcal{A}_Q = \mathbb{Z}$ -subalgebra of $\mathbb{Q}(x_1, \dots, x_N)$ gen. by \forall cluster variables.

Mutation at k : $\mu_k(Q, x) = (R, y)$

A new quiver R is obtained from Q by reversing \forall arrows incident with k and




\xrightarrow{r} means an $|r|$ -fold arrows $\begin{cases} \longrightarrow & (r \geq 0) \\ \longleftarrow & (r < 0) \end{cases}$

New cluster variables $y = (y_1, \dots, y_n)$ are given by

$$y_i = \begin{cases} x_i & i \neq k, \\ \frac{1}{x_k} \left(\prod_{\text{arrows } j \rightarrow k \text{ of } Q} x_j + \prod_{\text{arrows } k \rightarrow j \text{ of } Q} x_j \right) & i = k, \end{cases}$$

Example. $I = \{1, 2\}$. Initial seed $(Q, x) = (1 \rightarrow 2, \{a, b\})$.

Seeds denoted by $(a \rightarrow b)$, and mutation μ_1, μ_2 by --- .



Fomin-Zelevinsky theorem (2003) $\left\{ \begin{array}{l} (1) \text{ Laurent phenomenon,} \\ (2) \text{ Finite type classification.} \end{array} \right.$

(1) \forall cluster variables are Laurent polynomials.

(2) $\#\{\text{cluster var.}\} < \infty \Leftrightarrow Q^{\text{mutations}} \sim \text{orientation on ADE Dynkin diag.}$

Cluster algebra formulation of T-system. ($A_2, 4$) example

$$T_1^{(1)}(u-1)T_1^{(1)}(u+1) = T_2^{(1)}(u) + T_1^{(2)}(u),$$

$$T_2^{(2)}(u-1)T_2^{(2)}(u+1) = T_1^{(2)}(u)T_3^{(2)}(u) + T_2^{(1)}(u),$$

$$T_3^{(1)}(u-1)T_3^{(1)}(u+1) = T_2^{(1)}(u) + T_3^{(2)}(u).$$

$$\begin{array}{ccc}
 x_1 \longrightarrow x_4 & & T_1^{(1)}(0) \longrightarrow T_1^{(2)}(1) \\
 \uparrow \quad \downarrow & & \uparrow \quad \downarrow \\
 x_2 \longleftarrow x_5 & := & T_2^{(1)}(1) \longleftarrow T_2^{(2)}(0) \\
 \downarrow \quad \uparrow & & \downarrow \quad \uparrow \\
 x_3 \longrightarrow x_6 & & T_3^{(1)}(0) \longrightarrow T_3^{(2)}(1)
 \end{array}
 \xRightarrow{\mu_1\mu_3}
 \begin{array}{ccc}
 T_1^{(1)}(2) \longleftarrow T_1^{(2)}(1) & & \\
 \downarrow \quad \nearrow \quad \downarrow & & \\
 T_2^{(1)}(1) \longleftarrow T_2^{(2)}(0) & & \\
 \uparrow \quad \searrow \quad \uparrow & & \\
 T_3^{(1)}(2) \longleftarrow T_3^{(2)}(1) & & \\
 & & \mu_5 \downarrow
 \end{array}$$

$$\begin{array}{ccc}
 T_1^{(1)}(2) \longrightarrow T_1^{(2)}(3) & & T_1^{(1)}(2) \longrightarrow T_1^{(2)}(3) \\
 \uparrow \quad \downarrow & & \downarrow \quad \nwarrow \quad \downarrow \\
 T_2^{(1)}(3) \longleftarrow T_2^{(2)}(2) & \xleftarrow{\mu_2} & T_2^{(1)}(1) \longrightarrow T_2^{(2)}(2) \\
 \downarrow \quad \uparrow & & \uparrow \quad \nearrow \quad \uparrow \\
 T_3^{(1)}(2) \longrightarrow T_3^{(2)}(3) & & T_3^{(1)}(2) \longrightarrow T_3^{(2)}(3)
 \end{array}
 \xleftarrow{\mu_4\mu_6}
 \begin{array}{ccc}
 T_1^{(1)}(2) \longleftarrow T_1^{(2)}(1) & & \\
 \downarrow \quad \uparrow & & \\
 T_2^{(1)}(1) \longrightarrow T_2^{(2)}(2) & & \\
 \uparrow \quad \downarrow & & \\
 T_3^{(1)}(2) \longleftarrow T_3^{(2)}(1) & &
 \end{array}$$

$$(Q, x(u+2)) = \mu_2\mu_4\mu_6\mu_5\mu_3\mu_1(Q, x(u)) \quad \text{for } x = \{T_m^{(a)}\}.$$

Similarly, T-system for any (\mathfrak{g}, ℓ) is formulated as

$$(Q, x(u+2)) = \mu(Q, x(u)) \text{ by an appropriate choice of}$$

$$\begin{cases} Q & : \text{quiver,} \\ x(u) = \{x_i(u)\} & : \text{cluster variables suitably identified with } T_m^{(a)}(u)\text{'s,} \\ \mu = \mu_{i_1} \cdots \mu_{i_s} & : \text{composite mutation.} \end{cases}$$

(Full) periodicity for any (\mathfrak{g}, ℓ) is formulated as

$$(Q, x(u)) = \mu^{h^\vee + \ell}(Q, x(u)).$$

Y-system can be incorporated into [cluster algebra with coefficients](#).

Periodicity of T and Y-systems follows from periodicity in (generalized) cluster category.

Dilogarithm identity

$$L(x) = -\frac{1}{2} \int_0^x \left(\frac{\log(1-y)}{y} + \frac{\log y}{1-y} \right), \quad L(x) + L(1-x) = L(1) = \frac{\pi^2}{6}.$$

Let $Y_m^{(a)}(u) = Y_m^{(a)} > 0$ be the positive solution to the level ℓ restricted **constant** Y-system. (known to be unique)

Conjecture [A.N. Kirillov 1989]

$$\frac{6}{\pi^2} \sum_{a=1}^{\text{rank } \mathfrak{g}} t_a^{\ell-1} \sum_{m=1} L \left(\frac{Y_m^{(a)}}{1 + Y_m^{(a)}} \right) = \frac{\ell \dim \mathfrak{g}}{\ell + h^\vee} - \text{rank } \mathfrak{g} \quad (\ell \geq 1).$$

Example: $\mathfrak{g} = A_1$.

$$\frac{6}{\pi^2} \sum_{m=1}^{\ell-1} L \left(\frac{\sin^2 \frac{\pi}{\ell+2}}{\sin^2 \frac{\pi(m+1)}{\ell+2}} \right) = \frac{3\ell}{\ell+2} - 1.$$

Example: $\mathfrak{g} = A_n, D_n, E_n$. Constant Y-system reads

$$\sum_{m=1}^{\ell-1} C_{jm}^{(A_{\ell-1})} \log(1 + Y_m^{(a)})^{-1} = \sum_{b=1}^n C_{ab}^{(\mathfrak{g})} \log(1 + Y_j^{(b)}) \quad (C = \text{Cartan matrix}).$$

- utilized in so many calculations of central charges in
Integrable perturbations of CFT,
Low temperature specific heat & finite size corrections of spin chains,
Asymptotics of affine Lie algebra characters.
- proved in several individual cases by
Lewin, Watson, Kirillov, Richmond, Szekeres, Gliozzi, Tateo, Chapoton,...

Theorem. ([Nakanishi 2009](ADE), [IIKKN 2010](general \mathfrak{g}))
Dilogarithm conjecture is true for $\forall(\mathfrak{g}, \ell)$.

Idea of proof

- (1) Easier to prove more general functional dilogarithm identity

$$\frac{6}{\pi^2} \sum_{u \bmod \text{period}} \sum_{a=1}^{\text{rank } \mathfrak{g}} \sum_{m=1}^{t_a \ell - 1} L \left(\frac{Y_m^{(a)}(u)}{1 + Y_m^{(a)}(u)} \right) = \ell \dim \mathfrak{g} - (\ell + h^\vee) \text{rank } \mathfrak{g}.$$

- (2) LHS is invariant by varying $Y_m^{(a)}(u) > 0$ as long as Y-system is satisfied.
- (3) Letting $Y_k^{(b)}(v) \rightarrow 0$ for some (b, k, v) , the Y-system enforces all $Y_m^{(a)}(u)$'s tend to 0 or ∞ . (“Tropical Y-system”)
- (4) Count those $Y_m^{(a)}(u)$'s tending to ∞ and apply $L(1) = \frac{\pi^2}{6}$. □